

MISCELLANEOUS EXERCISE – 4

Q. 1 Multiple Choice Questions:

Four possible answers are given for the following questions. Tick (✓) the correct answer.

1. $(x+3)^2 = x^2 + 6x + 9$ is
 - (a) a linear equation
 - (b) an equation
 - (c) an identity
 - (d) none of these
2. $\frac{2x+1}{(x+1)(x-1)}$ is
 - (a) an improper fraction
 - (b) an equation
 - (c) a proper fraction
 - (d) none of these
3. $\frac{x^3+1}{(x-1)(x+2)}$ is
 - (a) a proper fraction
 - (b) an improper fraction
 - (c) an identity
 - (d) a constant term
4. A fraction in which the degree of numerator is less than the degree of the denominator is called
 - (a) an equation
 - (b) an improper fraction
 - (c) an identity
 - (d) a proper fraction
5. A function of the form $f(x) = \frac{N(x)}{D(x)}$, with $D(x) \neq 0$, where $N(x)$ and $D(x)$ are polynomials in x is called
 - (a) an identity
 - (b) an equation
 - (c) a fraction
 - (d) none of these
6. The identity $(5x+4)^2 = 25x^2 + 40x + 16$ is true for
 - (a) one value of x
 - (b) two values of x
 - (c) all values of x
 - (d) none of these

7. A fraction in which the degree of the numerator is greater or equal to the degree of denominator is called

- (a) a proper fraction
- (b) an improper fraction
- (c) an equation
- (d) algebraic relation

8. Partial fractions of $\frac{x-2}{(x-1)(x+2)}$ are

of the form

- (a) $\frac{A}{x-1} + \frac{B}{x+2}$
- (b) $\frac{Ax}{x-1} + \frac{B}{x+2}$
- (c) $\frac{A}{x-1} + \frac{Bx+C}{x+2}$
- (d) $\frac{Ax+B}{x-1} + \frac{C}{x+2}$

9. Partial fractions of $\frac{x+2}{(x+1)(x^2+2)}$ are

of the form

- (a) $\frac{A}{x+1} + \frac{B}{x^2+2}$
- (b) $\frac{A}{x+1} + \frac{Bx+C}{x^2+2}$
- (c) $\frac{Ax+B}{x+1} + \frac{C}{x^2+2}$
- (d) $\frac{A}{x+1} + \frac{Bx}{x^2+2}$

10. Partial fractions of $\frac{x^2+1}{(x+1)(x-1)}$ are

of the form

- (a) $\frac{A}{x+1} + \frac{B}{x-1}$
- (b) $1 + \frac{A}{x+1} + \frac{Bx+C}{x-1}$
- (c) $1 + \frac{A}{x+1} + \frac{B}{x-1}$
- (d) $\frac{Ax+B}{(x+1)} + \frac{C}{x-1}$

ANSWER KEY

1.	c	2.	c	3.	b	4.	d	5.	c
6.	c	7.	b	8.	a	9.	b	10.	c

Q. 2 Write short answers of the following questions:

(i) Define a rational fraction.

An expression of the form $\frac{N(x)}{D(x)}$ with $D(x) \neq 0$

and $N(x)$ and $D(x)$ are polynomials in x with real coefficients, is called a rational fraction. Every fractional expression can be expressed as a quotient of two polynomials.

(ii) What is a proper fraction?

A rational fraction $\frac{N(x)}{D(x)}$, with $D(x) \neq 0$ is

called a proper fraction if degree of the polynomial $N(x)$ in the numerator is less than the degree of the polynomial $D(x)$ in the denominator.

(iii) What is an improper fraction?

A rational fraction $\frac{N(x)}{D(x)}$, with $D(x) \neq 0$ is

called an improper fraction if degree of the polynomial $N(x)$ is greater or equal to the degree of the polynomial $D(x)$ e.g. $\frac{x^2+1}{x-1}$

(iv) What are partial fractions?

Every proper fraction $\frac{N(x)}{D(x)}$, with $D(x) \neq 0$

can be resolved into an algebraic sum of components fractions. These components fractions of a resultant fraction are called its partial fractions.

(v) How can we make partial fractions

of $\frac{x-2}{(x+2)(x+3)}$?

Solution: $\frac{x-2}{(x+2)(x+3)}$

Let $\frac{x-2}{(x+2)(x+3)} = \frac{A}{x+2} + \frac{B}{x+3} \dots\dots(i)$

Multiplying both sides by $(x+2)(x+3)$, we get
 $x-2 = A(x+3) + B(x+2) \dots\dots(ii)$

As both sides of the identity are equal for all values of x ,

Put $x+2=0$ i.e $x=-2$ in equation (ii), we get

$$-2-2 = A(-2+3)$$

$$-4 = A$$

$$\Rightarrow \boxed{A = -4}$$

Now put $x+3=0$ i.e $x=-3$ in equation (ii) we get

$$-3-2 = B(-3+2)$$

$$-5 = -B$$

$$\Rightarrow \boxed{B = 5}$$

Putting the value of A and B in equation(i) we get required partial fractions.

$$\frac{x-2}{(x+2)(x+3)} = \frac{-4}{x+2} + \frac{5}{x+3}$$

(vi) Resolve $\frac{1}{x^2-1}$ into partial fractions.

Solution: $\frac{1}{x^2-1} = \frac{1}{(x-1)(x+1)}$

Let $\frac{1}{(x-1)(x+1)} = \frac{A}{x-1} + \frac{B}{x+1} \dots\dots(i)$

Multiplying both sides By $(x-1)(x+1)$, we get

$$1 = A(x+1) + B(x-1) \dots\dots(ii)$$

As both sides of identity are equal for all values of x

Putting $x-1=0$ i.e $x=1$ in equation (ii) we get

$$1 = A(1+1)$$

$$1 = 2A$$

$$\Rightarrow \boxed{A = \frac{1}{2}}$$

Putting $x+1=0$ i.e $x=-1$ in equation (ii) we get

$$1 = B(-1-1)$$

$$1 = -2B$$

$$\Rightarrow \boxed{B = -\frac{1}{2}}$$

Putting the value of A and B in equation(i) we get required partial fractions.

$$\frac{1}{x^2-1} = \frac{1}{2(x-1)} - \frac{1}{2(x+1)}$$

(vii) Find partial fractions of $\frac{3}{(x+1)(x-1)}$

Solution: $\frac{3}{(x+1)(x-1)}$

$$\text{Let } \frac{3}{(x+1)(x-1)} = \frac{A}{x+1} + \frac{B}{x-1} \dots\dots\dots(i)$$

Multiplying both sides by $(x+1)(x-1)$, we get

$$3 = A(x-1) + B(x+1) \dots\dots\dots(ii)$$

As both sides of the identity are equal for all values of x .

Put $x+1=0$ i.e $x=-1$ put in equation (ii) we get

$$3 = A(-1-1)$$

$$3 = -2A \quad \Rightarrow \quad \boxed{A = -\frac{3}{2}}$$

Now put $x-1=0$ i.e $x=1$ in equation (ii) we get

$$\Rightarrow 3 = B(1+1)$$

$$3 = 2B \quad \Rightarrow \quad \boxed{B = \frac{3}{2}}$$

Putting the value of A and B in equation(i) we get required partial fractions.

$$\frac{3}{(x+1)(x-1)} = \frac{-3}{2(x+1)} + \frac{3}{2(x-1)} = \frac{3}{2} \left(\frac{1}{x-1} - \frac{1}{x+1} \right)$$

(viii) Resolve $\frac{x}{(x-3)^2}$ into partial fractions.

Solution: $\frac{x}{(x-3)^2}$

$$\text{Let } \frac{x}{(x-3)^2} = \frac{A}{x-3} + \frac{B}{(x-3)^2} \dots\dots\dots(i)$$

Multiplying both sides by $(x-3)^2$, we get

$$x = A(x-3) + B \dots\dots\dots(ii)$$

As both sides of the identity are equal for all values of x ,

Put $x-3=0$ i.e $x=3$ in equation (ii) we get

$$\Rightarrow \boxed{B = 3}$$

Now comparing the coefficients of x , we have

$$\Rightarrow \boxed{A = 1}$$

Putting the value of A and B in equation(i) we get required partial fractions.

$$\frac{x}{(x-3)^2} = \frac{1}{x-3} + \frac{3}{(x-3)^2}$$

(ix) How we can make the partial fractions of $\frac{x}{(x+a)(x-a)}$?

Solution: $\frac{x}{(x+a)(x-a)}$

$$\text{Let } \frac{x}{(x+a)(x-a)} = \frac{A}{x+a} + \frac{B}{x-a} \dots\dots\dots(i)$$

Multiplying both sides by $(x+a)(x-a)$, we get

$$x = A(x-a) + B(x+a) \dots\dots\dots(ii)$$

As both sides of the identity are equal for all values of x ,

Put $x+a=0$ i.e $x=-a$ put in equation (ii) we get

$$-a = A(-a-a)$$

$$-a = -2aA$$

$$\Rightarrow A = \frac{-a}{-2a}$$

$$\Rightarrow \boxed{A = \frac{1}{2}}$$

Now put $x-a=0$ i.e $x=a$ in equation (ii) we get

$$a = B(a+a)$$

$$a = 2aB$$

$$\Rightarrow B = \frac{a}{2a}$$

$$\Rightarrow \boxed{B = \frac{1}{2}}$$

Putting the value of A and B in equation(i) we get required partial fractions.

$$\begin{aligned} \frac{x}{(x+a)(x-a)} &= \frac{1}{2(x+a)} + \frac{1}{2(x-a)} \\ &= \frac{1}{2} \left(\frac{1}{x+a} + \frac{1}{x-a} \right) \end{aligned}$$

(x) Whether $(x+3)^2 = x^2 + 6x + 9$ is an identity?

Answer:

Yes $(x+3)^2 = x^2 + 6x + 9$ is an identity because it is true for all the values of x .