

EXERCISE 5.3

Q.1 If $U = \{1, 2, 3, 4, \dots, 10\}$
 $A = \{1, 3, 5, 7, 9\}$, $B = \{1, 4, 7, 10\}$
 then verify the following questions:

Solution:

(i) $A - B = A \cap B'$

$$\text{L.H.S} = A - B = \{1, 3, 5, 7, 9\} - \{1, 4, 7, 10\}$$

$$= \{3, 5, 9\} \dots\dots\dots (i)$$

$$\text{R.H.S} = A \cap B'$$

$$B' = U - B = \{1, 2, 3, \dots, 10\} - \{1, 4, 7, 10\}$$

$$= \{2, 3, 5, 6, 8, 9\}$$

$$A \cap B' = \{1, 3, 5, 7, 9\} \cap \{2, 3, 5, 6, 8, 9\}$$

$$= \{3, 5, 9\} \dots\dots\dots (ii)$$

From (i) and (ii)

$$\text{L.H.S} = \text{R.H.S}$$

$$A - B = A \cap B'$$

(ii) $B - A = B \cap A'$

$$\text{L.H.S} = B - A = \{1, 4, 7, 10\} - \{1, 3, 5, 7, 9\}$$

$$= \{4, 10\} \dots\dots\dots (i)$$

$$\text{R.H.S} = B \cap A'$$

$$A' = U - A = \{1, 2, 3, \dots, 10\} - \{1, 3, 5, 7, 9\}$$

$$= \{2, 4, 6, 8, 10\}$$

$$B \cap A' = \{1, 4, 7, 10\} \cap \{2, 4, 6, 8, 10\}$$

$$= \{4, 10\} \dots\dots\dots (ii)$$

From (i) and (ii)

$$\text{L.H.S} = \text{R.H.S}$$

$$B - A = B \cap A'$$

(iii) $(A \cup B)' = A' \cap B'$

$$\text{L.H.S} = (A \cup B)'$$

$$A \cup B = \{1, 3, 5, 7, 9\} \cup \{1, 4, 7, 10\}$$

$$= \{1, 3, 4, 5, 7, 9, 10\}$$

$$(A \cup B)' = U - (A \cup B)$$

$$= \{1, 2, 3, \dots, 10\} - \{1, 3, 4, 5, 7, 9, 10\}$$

$$= \{2, 6, 8\} \dots\dots\dots (i)$$

$$\text{R.H.S} = A' \cap B'$$

$$A' = U - A = \{1, 2, 3, \dots, 10\} - \{1, 3, 5, 7, 9\}$$

$$= \{2, 4, 6, 8, 10\}$$

$$B' = U - B = \{1, 2, 3, \dots, 10\} - \{1, 4, 7, 10\}$$

$$= \{2, 3, 5, 6, 8, 9\}$$

$$A' \cap B' = \{2, 4, 6, 8, 10\} \cap \{2, 3, 5, 6, 8, 9\}$$

$$= \{2, 6, 8\} \dots\dots\dots (ii)$$

From (i) and (ii)

$$\text{L.H.S} = \text{R.H.S}$$

(iv) $(A \cap B)' = A' \cup B'$

$$\text{L.H.S} = (A \cap B)'$$

$$A \cap B = \{1, 3, 5, 7, 9\} \cap \{1, 4, 7, 10\}$$

$$= \{1, 7\}$$

$$(A \cap B)' = U - (A \cap B)$$

$$= \{1, 2, 3, \dots, 10\} - \{1, 7\}$$

$$= \{2, 3, 4, 5, 6, 8, 9, 10\} \dots\dots (i)$$

$$\text{R.H.S} = A' \cup B'$$

$$A' = U - A = \{1, 2, 3, \dots, 10\} - \{1, 3, 5, 7, 9\}$$

$$= \{2, 4, 6, 8, 10\}$$

$$B' = U - B = \{1, 2, 3, \dots, 10\} - \{1, 4, 7, 10\}$$

$$= \{2, 3, 5, 6, 8, 9\}$$

$$A' \cup B' = \{2, 4, 6, 8, 10\} \cup \{2, 3, 5, 6, 8, 9\}$$

$$= \{2, 3, 4, 5, 6, 8, 9, 10\} \dots\dots (ii)$$

From (i) and (ii)

$$\text{L.H.S} = \text{R.H.S}$$

(v) $(A - B)' = A' \cup B$

$$\text{L.H.S} = (A - B)'$$

$$A - B = \{1, 3, 5, 7, 9\} - \{1, 4, 7, 10\}$$

$$= \{3, 5, 9\}$$

$$(A - B)' = U - (A - B)$$

$$= \{1, 2, 3, \dots, 10\} - \{3, 5, 9\}$$

$$= \{1, 2, 4, 6, 7, 8, 10\} \dots\dots (i)$$

$$\text{R.H.S} = A' \cup B$$

$$A' = U - A = \{1, 2, 3, \dots, 10\} - \{1, 3, 5, 7, 9\}$$

$$= \{2, 4, 6, 8, 10\}$$

$$A' \cup B = \{2, 4, 6, 8, 10\} \cup \{1, 4, 7, 10\}$$

$$= \{1, 2, 4, 6, 7, 8, 10\} \dots\dots (ii)$$

From (i) and (ii)

$$\text{L.H.S} = \text{R.H.S}$$

(vi) $(B - A)' = B' \cup A$

$$\text{L.H.S} = (B - A)'$$

$$B - A = \{1, 4, 7, 10\} - \{1, 3, 5, 7, 9\}$$

$$= \{4, 10\}$$

$$(B - A)' = U - (B - A)$$

$$= \{1, 2, 3, \dots, 10\} - \{4, 10\}$$

$$= \{1, 2, 3, 5, 6, 7, 8, 9\} \dots\dots (i)$$

$$\text{R.H.S} = B' \cup A$$

$$B' = U - B = \{1, 2, 3, 4, \dots, 10\} - \{1, 4, 7, 10\}$$

$$= \{2, 3, 5, 6, 8, 9\}$$

$$B' \cup A = \{2, 3, 5, 6, 8, 9\} \cup \{1, 3, 5, 7, 9\}$$

$$= \{1, 2, 3, 5, 6, 7, 8, 9\} \dots\dots (ii)$$

From (i) and (ii)

$$\text{L.H.S} = \text{R.H.S}$$

Q.2 If $A = \{1, 3, 5, 7, 9\}$ $B = \{1, 4, 7, 10\}$
 $C = \{1, 5, 8, 10\}$

Then verify the following:

Solution:

(i) $(A \cup B) \cup C = A \cup (B \cup C)$

L.H.S = $(A \cup B) \cup C$

= $(\{1, 3, 5, 7, 9\} \cup \{1, 4, 7, 10\}) \cup \{1, 5, 8, 10\}$

= $\{1, 3, 4, 5, 7, 9, 10\} \cup \{1, 5, 8, 10\}$

= $\{1, 3, 4, 5, 7, 8, 9, 10\}$ (i)

R.H.S = $A \cup (B \cup C)$

= $\{1, 3, 5, 7, 9\} \cup (\{1, 4, 7, 10\} \cup \{1, 5, 8, 10\})$

= $\{1, 3, 5, 7, 9\} \cup \{1, 4, 5, 7, 8, 10\}$

= $\{1, 3, 4, 5, 7, 8, 9, 10\}$ (ii)

From (i) and (ii)

L.H.S = R.H.S

(ii) $(A \cap B) \cap C = A \cap (B \cap C)$

L.H.S = $(A \cap B) \cap C$

= $(\{1, 3, 5, 7, 9\} \cap \{1, 4, 7, 10\}) \cap \{1, 5, 8, 10\}$

= $\{1, 7\} \cap \{1, 5, 8, 10\}$

= $\{1\}$ (i)

R.H.S = $A \cap (B \cap C)$

= $\{1, 3, 5, 7, 9\} \cap (\{1, 4, 7, 10\} \cap \{1, 5, 8, 10\})$

= $\{1, 3, 5, 7, 9\} \cap \{1, 10\}$

= $\{1\}$ (ii)

From (i) and (ii)

L.H.S = R.H.S

(iii) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

L.H.S = $A \cup (B \cap C)$

= $\{1, 3, 5, 7, 9\} \cup (\{1, 4, 7, 10\} \cap \{1, 5, 8, 10\})$

= $\{1, 3, 5, 7, 9\} \cup \{1, 10\}$

= $\{1, 3, 5, 7, 9, 10\}$ (i)

R.H.S = $(A \cup B) \cap (A \cup C)$

$A \cup B = \{1, 3, 5, 7, 9\} \cup \{1, 4, 7, 10\}$

= $\{1, 3, 4, 5, 7, 9, 10\}$

$A \cup C = \{1, 3, 5, 7, 9\} \cup \{1, 5, 8, 10\}$

= $\{1, 3, 5, 7, 8, 9, 10\}$

Now $(A \cup B) \cap (A \cup C)$

= $\{1, 3, 4, 5, 7, 9, 10\} \cap \{1, 3, 5, 7, 8, 9, 10\}$

= $\{1, 3, 5, 7, 9, 10\}$ (ii)

From (i) and (ii)

L.H.S = R.H.S

(iv) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

L.H.S = $A \cap (B \cup C)$

= $\{1, 3, 5, 7, 9\} \cap (\{1, 4, 7, 10\} \cup \{1, 5, 8, 10\})$

= $\{1, 3, 5, 7, 9\} \cap \{1, 4, 5, 7, 8, 10\}$

= $\{1, 5, 7\}$ (i)

R.H.S = $(A \cap B) \cup (A \cap C)$

$A \cap B = \{1, 3, 5, 7, 9\} \cap \{1, 4, 7, 10\}$

= $\{1, 7\}$

$A \cap C = \{1, 3, 5, 7, 9\} \cap \{1, 5, 8, 10\}$

= $\{1, 5\}$

Now $(A \cap B) \cup (A \cap C) = \{1, 7\} \cup \{1, 5\}$

= $\{1, 5, 7\}$ (ii)

From (i) and (ii)

L.H.S = R.H.S

Q.3 If $U = N$, then verify De-Morgan's laws by using:

$A = \phi, B = P$

Solution:

$A = \{ \}$

$B = \{2, 3, 5, 7, \dots\}$

$U = \{1, 2, 3, 4, 5, 6, 7, \dots\}$

(i) $(A \cup B)' = A' \cap B'$

L.H.S = $(A \cup B)'$

$A \cup B = \{ \} \cup \{2, 3, 5, 7, \dots\}$

= $\{2, 3, 5, 7, \dots\}$

$(A \cup B)' = U - (A \cup B)$

= $\{1, 2, 3, 4, 5, 6, 7, \dots\} - \{2, 3, 5, 7, \dots\}$

= $\{1, 4, 6, \dots\}$ (i)

R.H.S = $A' \cap B'$

$A' = U - A = \{1, 2, 3, 4, 5, 6, 7, \dots\} - \{ \}$

= $\{1, 2, 3, 4, 5, 6, 7, \dots\}$

$B' = U - B = \{1, 2, 3, 4, 5, 6, 7, \dots\} - \{2, 3, 5, 7, \dots\}$

= $\{1, 4, 6, \dots\}$

$A' \cap B' = \{1, 2, 3, 4, 5, 6, 7, \dots\} \cap \{1, 4, 6, \dots\}$

= $\{1, 4, 6, \dots\}$ (ii)

From (i) and (ii)

L.H.S = R.H.S

$$(ii) (A \cap B)' = A' \cup B'$$

$$L.H.S = (A \cap B)'$$

$$(A \cap B) = \{ \} \cap \{2, 3, 5, 7, \dots\}$$

$$= \{ \}$$

$$(A \cap B)' = U - (A \cap B)$$

$$= \{1, 2, 3, 4, 5, 6, 7, \dots\} - \{ \}$$

$$= \{1, 2, 3, 4, 5, 6, 7, \dots\} \dots \dots (i)$$

$$R.H.S = A' \cup B'$$

$$A' = U - A = \{1, 2, 3, 4, \dots\} - \{ \}$$

$$= \{1, 2, 3, 4, \dots\}$$

$$B' = U - B = \{1, 2, 3, 4, 5, 6, 7, \dots\} - \{2, 3, 5, 7, \dots\}$$

$$= \{1, 4, 6, \dots\}$$

$$A' \cup B' = \{1, 2, 3, 4, 5, 6, 7, \dots\} \cup \{1, 4, 6, \dots\}$$

$$= \{1, 2, 3, 4, \dots\} \dots \dots (ii)$$

From (i) and (ii)

$$L.H.S = R.H.S$$

Q.4 If $U = \{1, 2, 3, 4, \dots, 10\}$, $A = \{1, 3, 5, 7, 9\}$ and $B = \{2, 3, 4, 5, 8\}$ then prove the following questions by Venn Diagram.

Solution: $A \cap B = \{1, 3, 5, 7, 9\} \cap \{2, 3, 4, 5, 8\} = \{3, 5\}$

So given sets A and B are overlapping sets

$$(i) A - B = A \cap B'$$




Fig: 1 $(A - B)$




Fig: 2 $A \cap B'$

- $A - B$ is shown by horizontal line segments in fig. 1.
- B' is shown by vertical line segments and squares in fig. 2.
- $A \cap B'$ is shown by squares in fig. 2.

Regions shown in fig. 1 and fig. 2 are equal, thus $A - B = A \cap B'$

$$(ii) B - A = B \cap A'$$




Fig: 1 $(B - A)$




Fig: 2 $(B \cap A')$

- $B - A$ is shown by horizontal line segments in fig. 1.
- A' is shown by vertical line segments and squares in fig. 2.
- $B \cap A'$ is shown by squares in fig. 2.

Regions shown in fig. 1 and fig. 2 are equal, thus $B - A = B \cap A'$.

(iii) $(A \cup B)' = A' \cap B'$ (De-Morgan's Law)




Fig.1




Fig.2

- $A \cup B$ is shown by horizontal line segments in Fig. 1.
- $(A \cup B)'$ is shown by shaded area in Fig. 1.
- A' is shown by horizontal line segments and squares in Fig. 2.
- B' is shown by vertical line segments and squares in Fig. 2.
- $A' \cap B'$ is shown by squares in Fig. 2.

Shaded area shown in Fig. 1 and square area shown in Fig. 2 are equal.

thus $(A \cup B)' = A' \cap B'$

(iv) $(A \cap B)' = A' \cup B'$




Fig. 1




Fig. 2

- $A \cap B$ is shown by horizontal line segments in Fig. 1.
- $(A \cap B)'$ is shown by shaded area in Fig. 1.
- A' is shown by horizontal line segments and squares in Fig. 2.
- B' is shown by vertical line segments and squares in Fig. 2.
- $A' \cup B'$ is shown by squares, horizontal and vertical line segments in Fig. 2.

Shaded area shown in Fig. 1 and area of squares, vertical and horizontal line segments shown in Fig. 2 are equal. thus $(A \cap B)' = A' \cup B'$

(v) $(A - B)' = A' \cup B$




Fig. 1




Fig.2

- $A - B$ is shown by horizontal line segments in Fig. 1.
- $(A - B)'$ is shown by shaded area in Fig.1.
- A' is shown by squares Fig. 2.
- $A' \cup B$ is shown by squares and horizontal line segments in Fig. 2.

Shaded area in Fig. 1 and area of squares & horizontal line segments in Fig. 2 are equal.

thus $(A - B)' = A' \cup B$

$$(vi) \quad (B-A)' = B' \cup A$$




Fig. 1




Fig. 2

- $B-A$ is shown by horizontal line segments in Fig. 1.
 - $(B-A)'$ is shown by shaded area in Fig. 1.
 - B' is shown by squares in Fig. 2.
 - $B' \cup A$ is shown by squares and horizontal line segments in Fig. 2.
- Shaded area in Fig. 1 and area of squares & vertical line segments in Fig. 2 are equal.
thus $(B-A)' = B' \cup A$

Ordered Pairs and Cartesian Product:

(a) Ordered pairs

Any two numbers x and y , written in the form (x, y) is called an ordered pair. In an ordered pair (x, y) the order of numbers is important, that is, x is the first co-ordinate and y is the second co-ordinate. For example, $(3, 2)$ is different from $(2, 3)$.

It is obvious that $(x, y) \neq (y, x)$ unless $x = y$.

Note that $(x, y) = (s, t)$, iff $x = s$ and $y = t$

(b) Cartesian product:

Cartesian product of two non-empty sets A and B denoted by $A \times B$ consists of all ordered pairs (x, y) such that $x \in A$ and $y \in B$.

$$A \times B = \{(x, y) \mid x \in A \wedge y \in B\}$$

Example: If $A = \{1, 2, 3\}$ and $B = \{2, 5\}$, then find $A \times B$ and $B \times A$

Solution:

$$A \times B = \{1, 2, 3\} \times \{2, 5\}$$

$$A \times B = \{(1, 2), (1, 5), (2, 2), (2, 5), (3, 2), (3, 5)\}$$

Since set A has 3 elements and set B has 2 elements.

Hence product set $A \times B$ has $3 \times 2 = 6$ ordered pairs. We note that

$$B \times A = \{(2, 1), (2, 2), (2, 3), (5, 1), (5, 2), (5, 3)\}$$

Evidently $A \times B \neq B \times A$